

WITT VECTORS AND TAMBARA FUNCTORS

M. BRUN

ABSTRACT. We give, for every finite group G , a combinatorial description of the ring of G -Witt vectors on a polynomial algebra over the integers. Using this description we show that the functor, which takes a commutative ring with trivial action of G to its ring of Witt vectors, coincides with the left adjoint of the algebraic functor from the category of G -Tambara functors to the category of commutative rings with an action of G .

INTRODUCTION

In [Wi] Witt constructed an endofunctor on the category of commutative rings, which takes a commutative ring R to the ring $\mathbb{W}_p(R)$ of p -typical Witt vectors. This construction can be used to construct field extensions of the p -adic numbers [SW], and it is essential in the construction of crystalline cohomology [Be]. In [DS] Dress and Siebeneicher constructed for every pro-finite group G an endofunctor $\mathbb{W}_G$ on the category of commutative rings. In the case where $G = \widehat{C}_p$ is the pro- p -completion of the infinite cyclic group the functors $\mathbb{W}_p$ and $\mathbb{W}_G$ agree. The functor $\mathbb{W}_G$ is constructed in such a way that $\mathbb{W}_G(\mathbb{Z})$ is an appropriately completed Burnside ring for the pro-finite group G . For an arbitrary commutative ring R , the ring $\mathbb{W}_G(R)$ is somewhat mysterious, even when $G = \widehat{C}_p$. The first aim of the present paper is to give a new description of the ring $\mathbb{W}_G(R)$ when R is a polynomial algebra over the integers. In the special case where G is finite our description is given in terms of a category U^G introduced by Tambara in [Ta]. The category U^G has as objects the class of finite G -sets, and the morphism set $U^G(X, Y)$ is constructed as follows: Let $U_+^G(X, Y)$ denote the set of equivalence classes of chains of G -maps of the form $X \leftarrow A \rightarrow B \rightarrow Y$, where two chains $X \leftarrow A \rightarrow B \rightarrow Y$ and $X \leftarrow A' \rightarrow B' \rightarrow Y$ are defined to be equivalent if there exist G -bijections $A \rightarrow A'$ and $B \rightarrow B'$ making the following diagram commutative:

$$\begin{array}{ccccccc} X & \longleftarrow & A & \longrightarrow & B & \longrightarrow & Y \\ \parallel & & \downarrow & & \downarrow & & \parallel \\ X & \longleftarrow & A' & \longrightarrow & B' & \longrightarrow & Y. \end{array}$$

Date: April 30, 2003.

1991 Mathematics Subject Classification. 13K05; 18C10; 19A22.

Key words and phrases. Witt vectors, Mackey functors, Tambara functors, coloured theories.

The author wishes to thank the University of Osnabrück for hospitality and Rainer Vogt for a number of helpful conversations. This work has been supported by the DFG.

The set $U_+^G(X, Y)$ has a natural semi-ring-structure described in 1.2. By group-completing the underlying additive monoid of $U_+^G(X, Y)$ we obtain a commutative ring, and we define $U^G(X, Y)$ to be the set obtained by forgetting the ring-structure. For a subgroup H of G we let G/H denote the G -set of left cosets of H in G , and we let G/e denote G considered as a left G -set. The following theorem is a special case of theorem 5.5.

Theorem A. *Let G be a finite group and let X be a finite set considered as a G -set with trivial action. The ring $U^G(X, G/e)$ is the polynomial ring $\mathbb{Z}[X]$ over $\mathbb{Z}$, with one indeterminate for each element in X , and the ring $U^G(X, G/G)$ is naturally isomorphic to the ring $\mathbb{W}_G(U^G(X, G/e)) = \mathbb{W}_G(\mathbb{Z}[X])$.*

Our second aim is to advertise the category of G -Tambara functors, that is, the category $[U^G, \mathcal{E}ns]_0$ of set-valued functors on U^G preserving finite products. Tambara calls such a functor a TNR-functor, an acronym for “functor with trace, norm and restriction”, but we have chosen to call them G -Tambara functors. This category is intimately related to the Witt vectors of Dress and Siebeneicher. In order to explain this relation we consider the full subcategory U^{fG} of U^G with free G -sets as objects, and we note that the category of fG -Tambara functors, that is, the category $[U^{fG}, \mathcal{E}ns]_0$ of set-valued functors on U^{fG} preserving finite products is equivalent to the category of commutative rings with an action of G through ring-automorphisms. The categories U^G and U^{fG} are coloured theories in the sense of Boardman and Vogt [BV], and therefore they are complete and cocomplete, and the forgetful functor $[U^{fG}, \mathcal{E}ns]_0 \rightarrow [U^G, \mathcal{E}ns]_0$ induced by the inclusion $U^{fG} \subseteq U^G$ has a left adjoint functor, which we shall denote by L_G .

Theorem B. *Let G be a finite group. If R is a commutative ring with G acting trivially, then for every subgroup H of G , there is a natural isomorphism $\mathbb{W}_H(R) \cong L_G R(G/H)$.*

We are also able to describe the ring $L_G R(G/H)$ in the case where G acts nontrivially on R . In order to do this we need to introduce an ideal $\mathbb{I}_G(R)$ of $\mathbb{W}_G(R) = R^{\mathcal{Q}(G)}$, where $\mathcal{Q}(G)$ is the set of conjugacy classes of subgroups of G . An element $a \in \mathbb{W}_G(R)$ is of the form $a = (a_H)'_{H \leq G}$, where the prime means that $a_H = a_{gHg^{-1}} \in R$ for $H \leq G$ and $g \in G$. We let $\mathbb{I}_G(R) \subseteq \mathbb{W}_G(R)$ denote the ideal generated by elements of the form $a - b$, where $a = (a_K)'_{K \leq G}$ and $b = (b_K)'_{K \leq G}$ satisfy that for every $K \leq G$ there exist $n \geq 1$ and $g_{1,K}, \dots, g_{n,K} \in N_K(G)$ and $a_{1,K}, \dots, a_{n,K} \in R$ such that firstly $g_{1,K}K = \dots = g_{n,K}K$ and secondly $a_K = a_{1,K} \dots a_{n,K}$ and $b_K = (g_{1,K}a_{1,K}) \dots (g_{n,K}a_{n,K})$. Our description of $L_G R(G/H)$ is formulated in the following generalization of theorem B:

Theorem C. *Let G be a finite group and let R be a commutative ring with an action of G . There is a natural isomorphism $\mathbb{W}_H(R)/\mathbb{I}_H(R) \cong (L_G R)(G/H)$.*

Let us stress that in our setup the above result only makes sense for a finite group. For a pro-finite group G it might be possible to construct appropriate variants $\widehat{U}^G(X, Y)$ of the rings $U^G(X, Y)$ such that the ring $\widehat{U}^G(X, G/G)$ is naturally isomorphic to $\mathbb{W}_G(\mathbb{Z}[X])$. We shall not go further into this here.

There have been given other combinatorial constructions of the Witt vectors by Metropolis and Rota [MR] and Graham [Gra]. The ring of Witt vectors has been related to the ring of necklaces by Dress and Siebeneicher in [DS2]. Our combinatorial description differs from the previous ones mostly in that it is forced by some additional structure on the Witt vectors.

It is our belief that the category of G -Tambara functors has applications yet to be discovered. There is a rich supply of them coming from equivariant stable homotopy theory. In fact, every E_∞ ring G -spectrum gives rise to a G -Tambara functor by taking the zeroth homotopy group [Br]. In the case where $G = e$ is the trivial group, the category of G -Tambara functors is equivalent to the category of commutative rings. It is well known that every commutative ring can be realized as the zeroth homotopy group of an E_∞ ring-spectrum. For an arbitrary group G one may speculate whether G -Tambara functors can be realized as the zeroth homotopy group of an E_∞ ring-spectrum with an action of G .

Some concepts from commutative algebra can be carried over to the setting of G -Tambara functors. Most notably, the notions of an ideals, modules and chain complexes have G -Tambara versions. We leave these concepts to future work.

The paper is organized as follows: In section 1 we have collected some of the results from the papers [DS] and [Ta] that we need in the rest of the paper. In section 2 we note that the category of Tambara functors is the category of algebras for a coloured theory. In section 3 we construct a homomorphism relating Witt vectors and Tambara functors, which we have chosen to call the Teichmüller homomorphism. In section 4 we prove the fundamental fact that the Teichmüller homomorphism is a ring-homomorphism. In section 5 we prove that for free Tambara functors, the Teichmüller homomorphism is an isomorphism, and finally in section 6 we prove theorem C.

1. PREREQUISITES

In this section we fix some notation and recollect results from [DS] and [Ta]. All rings are supposed to be both commutative and unital. Given a group G we only consider left actions of G . A G -ring is a ring with an action of the group G through ring-automorphisms.

Given a pro-finite group G we let $\mathcal{O}(G)$ denote the G -set of open subgroups of G with action given by conjugation and we let $\underline{\mathcal{Q}}(G)$ denote the set of conjugacy classes of open subgroups of G . For a G -set X and a subgroup H of G we define $|X^H|$ to be the cardinality of the set X^H of H -invariant elements of X . The following is the main result of [DS].

Theorem 1.1. *Let G be a pro-finite group. There exists a unique endofunctor $\mathbb{W}_G$ on the category of rings such that for a ring R the ring $\mathbb{W}_G(R)$ has the set $R^{\underline{\mathcal{Q}}(G)}$ of maps from the set $\underline{\mathcal{Q}}(G)$ to R as underlying set, in such a way that for every ring-homomorphism $h : R \rightarrow R'$ and every $x \in \mathbb{W}_G(R)$ one has $\mathbb{W}_G(h)(x) = h \circ x$, while for any subgroup U of G the family of G -maps*

$$\phi_U^R : \mathbb{W}_G(R) \rightarrow R$$

defined by

$$x = (x_V)'_{V \leq G} \mapsto \sum'_{U \lesssim V \leq G} |U(G/V)^U| \cdot x_V^{(V:U)}$$

provides a natural transformation from the functor $\mathbb{W}_G$ into the identity functor. Here $U \lesssim V$ means that the subgroup U of G is subconjugate to V , i.e., there exists some $g \in G$ with $U \leq gVg^{-1}$, $(V : U)$ means the index of U in gVg^{-1} which coincides with $(G : U)/(G : V)$ and therefore is independent of g , and the symbol “ $\sum'$ ” is meant to indicate that for each conjugacy class of subgroups V with $U \lesssim V$ exactly one summand is taken, and an element $a \in \mathbb{W}_G(R)$ is written on the form $a = (a_V)'_{V \leq G}$, where the prime means that $a_V = a_{gVg^{-1}}$ for $g \in G$.

In the section 3 we shall reprove theorem 1.1.

The rest of this section is a recollection of the work [Ta] of Tambara. We let $\mathcal{F}in$ denote the category of finite sets and we let $\mathcal{F}in^G$ denote the category of finite G -sets. Given a finite G -set X we denote by $\mathcal{F}in^G/X$ the category of objects over X in $\mathcal{F}in^G$. Given $f : X \rightarrow Y$ in $\mathcal{F}in^G$ the pull-back functor

$$\begin{aligned} \mathcal{F}in^G/Y &\rightarrow \mathcal{F}in^G/X \\ (B \rightarrow Y) &\mapsto (X \times_Y B \rightarrow X) \end{aligned}$$

has a right adjoint

$$\begin{aligned} \Pi_f : \mathcal{F}in^U/X &\rightarrow \mathcal{F}in^G/Y \\ (A \xrightarrow{p} X) &\mapsto (\Pi_f A \xrightarrow{\Pi_f p} Y), \end{aligned}$$

where $\Pi_f p$ is made from p as follows. For each $y \in Y$, the fibre $(\Pi_f p)^{-1}(y)$ is the set of maps $s : f^{-1}(y) \rightarrow A$ such that $p(s(x)) = x$ for all $x \in f^{-1}(y)$. If $g \in G$ and $s \in (\Pi_f p)^{-1}(y)$, the map $^g s : f^{-1}(gy) \rightarrow A$ taking x to $gs(g^{-1}x)$ belongs to $(\Pi_f p)^{-1}(gy)$. The operation $(g, s) \mapsto ^g s$ makes $\Pi_f A$ a G -set and $\Pi_f p$ a G -map.

There is a commutative diagram

$$\begin{array}{ccccc} X & \xleftarrow{p} & A & \xleftarrow{e} & X \times_Y \Pi_f A \\ f \downarrow & & & & f' \downarrow \\ Y & \xleftarrow{\Pi_f p} & \Pi_f A & \xlongequal{\quad} & \Pi_f A, \end{array}$$

where f' is the projection and e is the evaluation map $(x, s) \mapsto s(x)$. A diagram in $\mathcal{F}in^G$ which is isomorphic to a diagram of the above form is called an *exponential diagram*.

We say that two diagrams $X \leftarrow A \rightarrow B \rightarrow Y$ and $X \leftarrow A' \rightarrow B' \rightarrow Y$ in $\mathcal{F}in^G$ are equivalent if there exist G -isomorphisms $A \rightarrow A'$, $B \rightarrow B'$ making the diagram

$$\begin{array}{ccccccc} X & \longleftarrow & A & \longrightarrow & B & \longrightarrow & Y \\ \parallel & & \downarrow & & \downarrow & & \parallel \\ X & \longleftarrow & A' & \longrightarrow & B' & \longrightarrow & Y \end{array}$$

commutative, and we let $U_+^G(X, Y)$ be the set of the equivalence classes $[X \leftarrow A \rightarrow B \rightarrow Y]$ of diagrams $X \leftarrow A \rightarrow B \rightarrow Y$.

Tambara defines an operation $\circ : U_+^G(Y, Z) \times U_+^G(X, Y) \rightarrow U_+^G(X, Z)$ by

$$[Y \leftarrow C \rightarrow D \rightarrow Z] \circ [X \leftarrow A \rightarrow B \rightarrow Y] = [X \leftarrow A'' \rightarrow \tilde{D} \rightarrow Z],$$

where the maps on the right are composites of the maps in the diagram

$$\begin{array}{ccccccc} X & \longleftarrow & A & \longleftarrow & A' & \longleftarrow & A'' \\ & & \downarrow & & \downarrow & & \downarrow \\ & & B & \longleftarrow & B' & \longleftarrow & \tilde{C} \\ & & \downarrow & & \downarrow & & \parallel \\ & & Y & \longleftarrow & C & & \tilde{C} \\ & & & & \downarrow & & \downarrow \\ & & & & Z & \longleftarrow & D & \longleftarrow & \tilde{D}. \end{array}$$

Here the three squares are pull-back diagrams and the diagram

$$\begin{array}{ccccc} C & \longleftarrow & B' & \longleftarrow & \tilde{C} \\ \downarrow & & & & \downarrow \\ D & \longleftarrow & \tilde{D} & \xlongequal{\quad} & \tilde{D} \end{array}$$

is an exponential diagram. He verifies that U_+^G is a category with $\circ$ as composition and given $f : X \rightarrow Y$ in $\mathcal{F}in^G$ he introduces the notation

$$\begin{aligned} R_f &= [Y \xleftarrow{f} X \rightrightarrows X \rightrightarrows X], \\ T_f &= [X \xleftarrow{\quad} X \rightrightarrows X \xrightarrow{f} Y] \quad \text{and} \\ N_f &= [X \xleftarrow{\quad} X \xrightarrow{f} Y \rightrightarrows Y]. \end{aligned}$$

Every morphism in U_+^G is a composition of morphisms on the above form. He also shows that every object X of U_+^G is a semi-ring object in the following natural way:

Proposition 1.2. *Given objects X and Y in U_+^G , there is semi-ring-structure on $U_+^G(X, Y)$ given as follows:*

$$\begin{aligned} 0 &= [X \leftarrow \emptyset \rightarrow \emptyset \rightarrow Y], \\ 1 &= [X \leftarrow \emptyset \rightarrow Y \rightarrow Y], \\ [X \leftarrow A \rightarrow B \rightarrow Y] + [X \leftarrow A' \rightarrow B' \rightarrow Y] &= [X \leftarrow A \amalg A' \rightarrow B \amalg B' \rightarrow Y] \end{aligned}$$

and

$$\begin{aligned} [X \leftarrow A \rightarrow B \rightarrow Y] \cdot [X \leftarrow A' \rightarrow B' \rightarrow Y] &= \\ [X \leftarrow B \times_Y A' \amalg A \times_Y B' \rightarrow B \times_Y B' \rightarrow Y]. \end{aligned}$$

It is also shown in [Ta] that there is a unique category U^G satisfying the following conditions:

- (i) $\text{ob } U^G = \text{ob } U_+^G$

- (ii) The morphism set $U^G(X, Y)$ is the group completion of the underlying additive monoid of $U_+^G(X, Y)$.
- (iii) The group completion maps $k : U_+^G(X, Y) \rightarrow U^G(X, Y)$ and the identity on $\text{ob}(U_+^G)$ form a functor $k : U_+^G \rightarrow U^G$.
- (iv) The functor k preserves finite products.

Proposition 1.3. (i) If $X_1 \xrightarrow{i_1} X \xleftarrow{i_2} X_2$ is a sum diagram in $\mathcal{F}in^G$, then $X_1 \xleftarrow{R_{i_1}} X \xrightarrow{R_{i_2}} X_2$ is a product diagram in U^G and $\emptyset$ is final in U^G .

(ii) Let X be a G -set and $\nabla : X \amalg X \rightarrow X$ the folding map, $i : \emptyset \rightarrow X$ the unique map. Then X has a structure of a ring object of U^G with addition T_∇ , additive unit T_i , multiplication N_∇ and multiplicative unit N_i .

(iii) If $f : X \rightarrow Y$ is a G -map, then the morphisms R_f , T_f and N_f of U^G preserve the above structures of ring, additive group and multiplicative monoid on X and Y , respectively.

Given a category $\mathcal{C}$ with finite products, we shall denote the category of set-valued functors on $\mathcal{C}$ preserving finite products by $[\mathcal{C}, \mathcal{E}ns]_0$. The morphisms in $[\mathcal{C}, \mathcal{E}ns]_0$ are given by natural transformations.

Definition 1.4. The category of G -Tambara functors is the category $[U^G, \mathcal{E}ns]_0$. That is, a G -Tambara functor (called TNR-functor in [Ta]) is a set-valued functor on U^G preserving finite products, and a morphism of G -Tambara functors is a natural transformation.

Given a G -Tambara functor S and $[X \leftarrow A \rightarrow B \rightarrow Y] \in U^G(X, Y)$ we obtain a function $S[X \leftarrow A \rightarrow B \rightarrow Y] : S(X) \rightarrow S(Y)$. Since S is product-preserving, it follows from (ii) of proposition 1.3 that $S(X)$ is a ring. Given a finite G -map $f : X \rightarrow Y$ we shall use the notation $S^*(f) = S(R_f)$, $S_+(f) = S(T_f)$ and $S_\bullet(f) = S(N_f)$. It follows from (iii) of 1.3 that $S^*(f)$ is a ring-homomorphism, that $S_+(f)$ is an additive homomorphism and that $S_\bullet(f)$ is multiplicative. A G -Tambara functor S is uniquely determined by the functions $S^*(f)$, $S_+(f)$ and $S_\bullet(f)$ for all $f : X \rightarrow Y$ in $\mathcal{F}in^G$.

Given subgroups $K \leq H \leq G$ we shall denote by $\pi_K^H : G/K \rightarrow G/H$ the projection induced by the inclusion $K \leq H$ and given $g \in G$ we shall let $c_g : G/H \rightarrow G/gHg^{-1}$ denote conjugation by g , $c_g(\sigma H) = \sigma g^{-1}(gHg^{-1})$.

2. COLOURED THEORIES

In this section we shall explain that the category U^G is an $\mathcal{Q}(G)$ -coloured category in the sense of Boardman and Vogt [BV].

- Definition 2.1** ([BV, 2.3,]). (i) Let $\mathcal{O}$ be a finite set. An $\mathcal{O}$ -coloured theory is a category Θ together with a faithful functor $\sigma_\Theta : (\mathcal{F}in/\mathcal{O})^{\text{op}} \rightarrow \Theta$ such that firstly σ_Θ preserves finite products and secondly every object of Θ is isomorphic to an object in the image of σ_Θ .
- (ii) The category of algebras over a theory Θ is the category $[\Theta, \mathcal{E}ns]_0$ of product-preserving set-valued functors on Θ .

(iii) A morphism $\gamma : \Theta \rightarrow \Psi$ of $\mathcal{O}$ -coloured theories is a functor preserving finite products.

Other authors, e.g. [AR] and [Re], use the name “sorted theory” for a coloured theory.

Given a finite group G , choosing representatives G/H for the objects of $\underline{\mathcal{Q}}(G)$, we can construct a functor

$$\begin{aligned} \sigma_{(\mathcal{F}in^G)^{\text{op}}} : (\mathcal{F}in/\underline{\mathcal{Q}}(G))^{\text{op}} &\rightarrow (\mathcal{F}in^G)^{\text{op}}, \\ (z : Z \rightarrow \underline{\mathcal{Q}}(G)) &\mapsto \coprod_{[G/H] \in \underline{\mathcal{Q}}(G)} G/H \times z^{-1}([G/H]). \end{aligned}$$

This way we give $(\mathcal{F}in^G)^{\text{op}}$ the structure of an $\underline{\mathcal{Q}}(G)$ -coloured theory. Composing $\sigma_{(\mathcal{F}in^G)^{\text{op}}}$ with the functor $R : (\mathcal{F}in^G)^{\text{op}} \rightarrow U^G$, $f \mapsto R_f$ we obtain by (i) of 1.3 a functor $\sigma_{U^G} : (\mathcal{F}in/\underline{\mathcal{Q}}(G))^{\text{op}} \rightarrow U^G$ preserving finite products, and we obtain a structure of $\underline{\mathcal{Q}}(G)$ -coloured theory on U^G . The functor R is a morphism of $\underline{\mathcal{Q}}(G)$ -coloured theories.

Let $V^G \subseteq U^G$ denote the subcategory of U^G with the same class of objects as U^G and with $V^G(X, Y) \subseteq U^G(X, Y)$ given by the subgroup generated by morphisms of the form $[X \leftarrow A \xrightarrow{\equiv} A \rightarrow Y]$. The inclusion $V^G \subseteq U^G$ preserves finite products, and we have morphisms $(\mathcal{F}in^G)^{\text{op}} \rightarrow V^G \rightarrow U^G$ of $\underline{\mathcal{Q}}(G)$ -coloured theories. The category V^G is strongly related to the category spans considered by Lindner in [Li], and in fact the category of Mackey functors is equal to the category of V^G -algebras. Let $\mathcal{F}in^{fG}$ denote the full subcategory of $\mathcal{F}in^G$ with finite free G -sets as objects. The functor

$$\sigma_{(\mathcal{F}in^{fG})^{\text{op}}} : (\mathcal{F}in/\underline{\mathcal{Q}}(G))^{\text{op}} \rightarrow (\mathcal{F}in^{fG})^{\text{op}}, \quad (z : Z \rightarrow \underline{\mathcal{Q}}(G)) \mapsto G/e \times z^{-1}([G/e])$$

gives $(\mathcal{F}in^{fG})^{\text{op}}$ the structure of an $\underline{\mathcal{Q}}(G)$ -coloured theory. Similarly we can consider the $\underline{\mathcal{Q}}(G)$ -coloured theories given by the full subcategories $U^{fG} \subseteq U^G$ and $V^{fG} \subseteq V^G$ with finite free G -sets as objects. We have the following diagram of morphisms of $\underline{\mathcal{Q}}(G)$ -coloured theories:

$$\begin{array}{ccccc} (\mathcal{F}in^{fG})^{\text{op}} & \longrightarrow & V^{fG} & \longrightarrow & U^{fG} \\ \downarrow & & \downarrow & & \downarrow \\ (\mathcal{F}in^G)^{\text{op}} & \longrightarrow & V^G & \longrightarrow & U^G, \end{array}$$

where the vertical functors are inclusions of full subcategories.

- Lemma 2.2.** (i) *The category $[U^{fG}, \mathcal{E}ns]_0$ of fG -Tambara functors is equivalent to the category of G -rings.*
(ii) *The category $[V^{fG}, \mathcal{E}ns]_0$ of V^{fG} -algebras is equivalent to the category of left $\mathbb{Z}[G]$ -modules.*
(iii) *The category $[(\mathcal{F}in^{fG})^{\text{op}}, \mathcal{E}ns]_0$ of $(\mathcal{F}in^{fG})^{\text{op}}$ -algebras is equivalent to the category of G -sets.*

Proof. Since the statements have similar proofs we only give the proof of (i). Given an fG -Tambara functor S , we construct a G -ring-structure on $R = S(G/e)$. Indeed

by (ii) of 1.3 $S(G/e)$ is a ring, and given $g \in G$ the right multiplication $g : G/e \rightarrow G/e$, $x \mapsto xg$, induces a ring-automorphism $S^*(g^{-1})$ of $R = S(G/e)$. From the functoriality of S we obtain that R is a G -ring. Conversely, given a G -ring R' we shall construct an fG -Tambara functor S' . We define $S'(X)$ to be the set of G -maps from X to R' . Given $[X \xleftarrow{d} A \xrightarrow{f} B \xrightarrow{g} Y] \in U^G(X, Y)$, we define $S'[X \xleftarrow{d} A \xrightarrow{f} B \xrightarrow{g} Y] : S'(X) \rightarrow S'(Y)$ by the formula

$$S'[X \xleftarrow{d} A \xrightarrow{f} B \xrightarrow{g} Y](\phi)(y) = \sum_{b \in g^{-1}(y)} \left(\prod_{a \in f^{-1}(b)} \phi(d(a)) \right)$$

for $\phi \in S'(X)$ and $y \in Y$. We leave it to the reader to check that $S \mapsto R$ and $R' \mapsto S'$ are inverse functors up to isomorphism. $\square$

We refer to [Re, Propositions 4.3 and 4.7.] for a proof of the following two results. Alternatively the reader may modify the proofs given in [Bo, 3.4.5 and 3.7.7] for their monochrome versions.

Proposition 2.3. *Let Θ be an $\mathcal{O}$ -coloured theory. The category of Θ -algebras is complete and cocomplete.*

Proposition 2.4. *Given a morphism $\gamma : \Theta \rightarrow \Psi$ of $\mathcal{O}$ -coloured theories, the functor $\gamma^* : [\Psi, \mathcal{E}ns]_0 \rightarrow [\Theta, \mathcal{E}ns]_0$, $A \mapsto A \circ \gamma$ has a left adjoint $\gamma_* : [\Theta, \mathcal{E}ns]_0 \rightarrow [\Psi, \mathcal{E}ns]_0$.*

Definition 2.5. The category of fG -Tambara functors is the category $[U^{fG}, \mathcal{E}ns]_0$ of U^{fG} -algebras.

Definition 2.6. We let $L_G = i_* : [U^{fG}, \mathcal{E}ns]_0 \rightarrow [U^G, \mathcal{E}ns]_0$ denote the left adjoint of the functor $i^* : [U^G, \mathcal{E}ns]_0 \rightarrow [U^{fG}, \mathcal{E}ns]_0$ induced by the inclusion $i : U^{fG} \subseteq U^G$.

Let us note that L_G can be constructed as the left Kan extension along i , and that for $R \in [U^{fG}, \mathcal{E}ns]_0$, we have for every finite free G -set X an isomorphism $L_G R(X) \cong R(X)$ because U^{fG} is a full subcategory of U^G .

3. THE TEICHMÜLLER HOMOMORPHISM

We shall now give a connection between the category of G -Tambara functors and the category of rings with an action of G . Recall that U^{fG} denotes the full subcategory of U^G with objects given by free G -sets. Below we use notation introduced in 1.1.

Definition 3.1. Let G be a finite group and let S be a G -Tambara functor. The *unrestricted Teichmüller homomorphism* $t : \mathbb{W}_G(S(G/e)) \rightarrow S(G/G)$ takes $(x_U)'_{U \leq G} \in \mathbb{W}_G(S(G/e))$ to

$$t((x_U)'_{U \leq G}) = \sum'_{U \leq G} S_+(\pi_U^G) S_\bullet(\pi_e^U)(x_U).$$

We shall prove the following proposition in the next section.

Proposition 3.2. *The unrestricted Teichmüller homomorphism $t : \mathbb{W}_G(S(G/e)) \rightarrow S(G/G)$ is a ring-homomorphism.*

In general t will neither be injective nor surjective. However, in the introduction, we have given an ideal $\mathbb{I}_G(S(G/e)) \subseteq \mathbb{W}_G(S(G/e))$ contained in the kernel of t :

Definition 3.3. Let G be a finite group and let R be a commutative ring with an action of G . We let $\mathbb{I}_G(R) \subseteq \mathbb{W}_G(R)$ denote the ideal generated by elements of the form $a - b$, where $a = (a_K)'_{K \leq G}$ and $b = (b_K)'_{K \leq G}$ satisfy that for every $K \leq G$ there exist $n \geq 1$ and $g_{1,K}, \dots, g_{n,K} \in N_K(G)$ and $a_{1,K}, \dots, a_{n,K} \in R$ such that firstly $g_{1,K}K = \dots = g_{n,K}K$ and secondly $a_K = a_{1,K} \dots a_{n,K}$ and $b_K = (g_{1,K}a_{1,K}) \dots (g_{n,K}a_{n,K})$.

Lemma 3.4. *Let S be a G -Tambara functor. The unrestricted Teichmüller homomorphism $t : \mathbb{W}_G(S(G/e)) \rightarrow S(G/G)$ maps the ideal $\mathbb{I}_G(S(G/e))$ to zero.*

Proof. It suffices to consider the case where $a_U = 0$ for $U \neq K$. We have that

$$\begin{aligned} & S_+(\pi_K^G)S_\bullet(\pi_e^K)(g_1a_1 \dots g_na_n) \\ &= S[\prod_1^n G/e \xleftarrow{\prod_{i=1}^n g_i} \prod_1^n G/e \xrightarrow{\pi_e^K \circ \nabla} G/K \xrightarrow{\pi_K^G} G/G](a_1, \dots, a_n) \\ &= S[\prod_1^n G/e \xleftarrow{\quad} \prod_1^n G/e \xrightarrow{\pi_e^K \circ \nabla} G/K \xrightarrow{\pi_K^G} G/G](a_1, \dots, a_n) \\ &= S_+(\pi_K^G)S_\bullet(\pi_e^K)(a_1 \dots a_n), \end{aligned}$$

where $\nabla : \prod_1^n G/e \rightarrow G/e$ is the fold map. $\square$

Definition 3.5. Let G be a finite group and let S be a G -Tambara functor. The Teichmüller homomorphism $\tau : \mathbb{W}_G(S(G/e))/\mathbb{I}_G(S(G/e)) \rightarrow S(G/G)$ is the ring-homomorphism induced by the unreduced Teichmüller homomorphism t .

The following theorem, proved in section 5, implies the case $H = G$ of theorem C.

Theorem 3.6. *Let G be a finite group and let R be an fG -Tambara functor. The Teichmüller homomorphism*

$$\tau : \mathbb{W}_G(L_GR(G/e))/\mathbb{I}_G(L_GR(G/e)) \rightarrow L_GR(G/G)$$

is an isomorphism. In particular, if G acts trivially on R , the Teichmüller homomorphism is an isomorphism on the form $\tau : \mathbb{W}_G(L_GR(G/e)) \rightarrow L_GR(G/G)$

In the above situation there is an isomorphism $L_GR(G/e) \cong R(G/e)$.

4. WITT POLYNOMIALS

We shall prove the following version of a theorem of Dress and Siebeneicher [DS, Theorem 3.2]. As we shall see below, theorem 1.1 and proposition 3.2 are immediate consequences of it.

Theorem 4.1. *Given a finite group G , there exist unique families $(s_U)'_{U \leq G}$, $(p_U)'_{U \leq G}$ of integral polynomials*

$$s_U = s_U^G, \quad p_U = p_U^G \in \mathbb{Z}[a_V, b_V \mid U \lesssim V \leq G]$$

in two times as many variables x_V, y_V ($U \lesssim V \leq G$) as there are conjugacy classes of subgroups $V \leq G$ which contain a conjugate of U such that for every G -Tambara functor S and for every $x = (x_U)'_{U \leq G}$ and $y = (y_U)'_{U \leq G} \in \mathbb{W}_G(S(G/e))$ we have that

$$\begin{aligned} \tau(x) + \tau(y) &= \tau((s_U(x_V, y_V | U \lesssim V \leq G))'_{U \leq G}) \quad \text{and} \\ \tau(x) \cdot \tau(y) &= \tau((p_U(x_V, y_V | U \lesssim V \leq G))'_{U \leq G}). \end{aligned}$$

Similarly, there exist polynomials $m_U = m_U^G \in \mathbb{Z}[a_V | U \lesssim V \leq G]$ such that for every G -Tambara functor S and for every $x = (x_U)'_{U \leq G} \in \mathbb{W}_G(S(G/e))$ we have that $-\tau(x) = \tau((m_U(x_V | U \lesssim V \leq G))'_{U \leq G})$. Further, for every subgroup H of G we have that

$$\begin{aligned} \phi_H(x) + \phi_H(y) &= \phi_H((s_U(x_V, y_V | U \lesssim V \leq G))'_{U \leq G}) \quad \text{and} \\ \phi_H(x) \cdot \phi_H(y) &= \phi_H((p_U(x_V, y_V | U \lesssim V \leq G))'_{U \leq G}). \end{aligned}$$

We shall call the polynomials s_U, p_U and m_U the Witt polynomials.

Proof of theorem 1.1. We first consider the case where G is finite. Given a ring R , we define operations $+$ and $\cdot$ on $\mathbb{W}_G(R)$ by defining $(a_U)'_{U \leq G} + (b_U)'_{U \leq G} := (s_U(a_V, b_V | U \lesssim V \leq G))'_{U \leq G}$ and $(a_U)'_{U \leq G} \cdot (b_U)'_{U \leq G} := (p_U(a_V, b_V | U \lesssim V \leq G))'_{U \leq G}$. In the case where R has no torsion, the map $\phi : \mathbb{W}_G(R) \rightarrow \prod'_{U \leq G} R$ with U 'th component ϕ_U is injective, and hence $\mathbb{W}_G(R)$ is a subring of $\prod'_{U \leq G} R$. In the case where R has torsion, we can choose a surjective ring-homomorphism $R' \rightarrow R$ from a torsion free ring R' . We obtain a surjection $\mathbb{W}_G(R') \rightarrow \mathbb{W}_G(R)$ respecting the operations $\cdot$ and $+$. Since $\mathbb{W}_G(R')$ is a ring we can conclude that $\mathbb{W}_G(R)$ is a ring. Given a surjective homomorphism $\gamma : G \rightarrow G'$ of finite groups we obtain a ring-homomorphism $\text{restr}_{G'}^G : \mathbb{W}_G(R) \rightarrow \mathbb{W}_{G'}(R)$ with $\text{restr}_{G'}^G((a_U)'_{U \leq G}) = ((b_V)')_{V \leq G'}$, where $b_V = a_{\gamma^{-1}(V)}$. (See [DS, (3.3.11)].) The easiest way to see that $\text{restr}_{G'}^G$ is a ring-homomorphism is to note that $(\gamma^{-1}H : \gamma^{-1}U) = (H : U)$ and that $\phi_{\gamma^{-1}H}(G/\gamma^{-1}U) = \phi_H(G'/H)$. For the case where G is a pro-finite group we note that for the ring $\mathbb{W}_G(R)$ can be defined to be the limit $\lim_N \mathbb{W}_{G/N}(R)$ taken over all finite factor groups G/N , with respect maps on the form $\text{restr}_{G'}^G$. $\square$

Proof of proposition 3.2. Proposition 3.2 follows from the first part of theorem 4.1 because we use the Witt polynomials to define the ring-structure on the Witt vectors. $\square$

For the rest of this section we fix a finite group G . We shall follow [DS] closely in the proof of theorem 4.1. For the uniqueness of the Witt polynomials we consider the representable G -Tambara functor $\Omega := U^G(\emptyset, -)$. In [DS, Theorem 2.12.7] it is shown that $\tau : \prod'_{U \leq G} \mathbb{Z} = \mathbb{W}_G(\Omega(G/e)) \rightarrow \Omega_G(\mathbb{Z})$ is a bijection. Hence the Witt polynomials are unique. We shall use the following six lemmas to establish the existence of Witt polynomials with the properties required in theorem 4.1.

Lemma 4.2. *For a subset A of G , let $U_A := \{g \in G \mid Ag = A\}$ denote its stabilizer group and let $i_A := |A/U_A|$ denote the number of U_A -orbits in A . If the set $\mathcal{U}(G)$ of subsets of G is considered as a G -set via $G \times \mathcal{U}(G) \rightarrow \mathcal{U}(G) : (g, A) \mapsto Ag^{-1}$, then*

for any $s, t \in S(G/e)$ one has

$$S_{\bullet}(\pi_e^G)(s+t) = \sum_{G \cdot A \in G \setminus \mathcal{U}(G)} S_+(\pi_{U_A}^G) S_{\bullet}(\pi_e^{U_A})(s^{i_A} \cdot t^{i_{G-A}}).$$

Proof. We let $i_1, i_2 : G/e \rightarrow G/e \amalg G/e$ denote the two natural inclusions. We have an exponential diagram

$$\begin{array}{ccccc} G/e & \xleftarrow{\nabla} & G/e \amalg G/e & \xleftarrow{d} & G/e \times \mathcal{U}(G) \\ \pi_e^G \downarrow & & & & \text{pr} \downarrow \\ G/G & \xleftarrow{\quad} & \mathcal{U}(G) & \xlongequal{\quad} & \mathcal{U}(G), \end{array}$$

where $d(g, A) = i_1(g)$ if $g^{-1} \in A$ and $d(g, A) = i_2(g)$ if $g^{-1} \notin A$. Since $\mathcal{U}(G) \cong \coprod_{G \cdot A \in G \setminus \mathcal{U}(G)} G \cdot A$, we have that

$$\begin{aligned} S_{\bullet}(\pi_e^G)(s+t) &= S_{\bullet}(\pi_e^G) S_+(\nabla)(s, t) \\ &= S[G/e \amalg G/e \xleftarrow{d} G/e \times \mathcal{U}(G) \rightarrow \mathcal{U}(G) \rightarrow G/G](s, t) \\ &= \sum_{G \cdot A \in G \setminus \mathcal{U}(G)} S[G/e \amalg G/e \xleftarrow{d} G/e \times GA \rightarrow GA \rightarrow G/G](s, t) \\ &= \sum_{G \cdot A \in G \setminus \mathcal{U}(G)} S[G/e \amalg G/e \leftarrow G/e \times A/U_A \amalg G/e \times (G-A)/U_A \rightarrow G/U_A \rightarrow G/G](s, t) \\ &= \sum_{G \cdot A \in G \setminus \mathcal{U}(G)} S[G/e \xleftarrow{\quad} G/e \xrightarrow{\pi_e^{U_A}} G/U_A \xrightarrow{\pi_{U_A}^G} G/G](s^{i_A} t^{i_{G-A}}) \\ &= \sum_{G \cdot A \in G \setminus \mathcal{U}(G)} S_+(\pi_{U_A}^G) S_{\bullet}(\pi_e^{U_A})(s^{i_A} t^{i_{G-A}}), \end{aligned}$$

where the maps without labels are natural projections. $\square$

Lemma 4.3. *Let R be a commutative ring. With the notation of lemma 4.2, we have for every subset A and subgroup U of G and for every $s, t \in R$, that*

$$(s+t)^{(G:U)} = \sum_{G \cdot A \in G \setminus \mathcal{U}(G)} |U(G/U_A)^U| \cdot (s^{i_A} \cdot t^{i_{G-A}})^{(U_A:U)}.$$

Proof. We compute:

$$\begin{aligned} (s+t)^{(G:U)} &= \sum_{A \subseteq G/U} s^{|A|} t^{|G/U|-|A|} = \sum_{A \in \mathcal{U}(G), U \leq U_A} s^{|A/U|} t^{|(G-A)/U|} \\ &= \sum_{G \cdot A \in G \setminus \mathcal{U}(G)} |(G/U_A)^U| \cdot (s^{i_A} \cdot t^{i_{G-A}})^{(U_A:U)}. \end{aligned}$$

$\square$

The following lemma is a variation on [DS, lemma 3.2.5], and the proof we give here is essentially identical to the one given in [DS].

Lemma 4.4. *For some $k \in \mathbb{N}$ let $V_1, \dots, V_k \leq G$ be a sequence of subgroups of G . Then for every subgroup $U \leq G$ and every sequence $\varepsilon_1, \dots, \varepsilon_k \in \{\pm 1\}$ there exists a polynomial $\xi_U = \xi_{(U; V_1, \dots, V_k; \varepsilon_1, \dots, \varepsilon_k)}^G = \xi_U(x_1, \dots, x_k) \in \mathbb{Z}[x_1, \dots, x_k]$ such that for every G -Tambara functor S and all $s_1, \dots, s_k \in S(G/e)$ one has*

$$\sum_{i=1}^k \varepsilon_i S_+(\pi_{V_i}^G) S_\bullet(\pi_e^{V_i})(s_i) = \sum'_{U \leq G} S_+(\pi_U^G) S_\bullet(\pi_e^U)(\xi_U(s_1, \dots, s_k)).$$

Proof. If $\varepsilon_1 = \varepsilon_2 = \dots = \varepsilon_k = 1$ and if V_i is not conjugate to V_j for $i \neq j$, then

$$\sum_{i=1}^k \varepsilon_i S_+(\pi_{V_i}^G) S_\bullet(\pi_e^{V_i})(s_i) = \sum'_{U \leq G} S_+(\pi_U^G) S_\bullet(S_e^U)(s_U),$$

with $s_U = s_i$ if U is conjugate to V_i and $s_U = 0$ if U is not conjugate to any of the $V_1, \dots, V_k$. So in this case we are done: $\xi_U(s_1, \dots, s_k) = s_U$. We use triple induction to prove the lemma. First with respect to $m_1 = m_1(U; V_1, \dots, V_k; \varepsilon_1, \dots, \varepsilon_k) := \max(|V_i| \mid \varepsilon_i = -1 \text{ or there exists some } j \neq i \text{ with } V_j \sim V_i)$, then with respect to $m_3 := |\{i \mid |V_i| = m_1 \text{ and there exists some } j \neq i \text{ with } V_j \sim V_i\}|$ and then with respect to $m_2 := |\{i \mid |V_i| = m_1 \text{ and } \varepsilon_i = -1\}|$.

We have just verified that the lemma holds in the case $m_1 = 0$. In case $m_1 > 0$ we have either $m_2 > 0$ or $m_3 > 0$. In case $m_2 > 0$, say $|V_1| = m_1$ and $\varepsilon_1 = -1$, we may use 4.2 with $G = V_1$, $s = -s_1$, $t = s_1$ to conclude that

$$0 = \sum_{V_1 \cdot A \in V_1 \backslash \mathcal{U}(V_1)} S_+(\pi_{U_A}^{V_1}) S_\bullet(\pi_e^{U_A})((-1)^{i_A} s_1^{(V_1:U_A)}).$$

Therefore, considering the two special summands $A = \emptyset$ and $A = V_1$ and putting $\mathcal{U}_0(V_1) := \{A \in \mathcal{U}(V_1) \mid A \neq \emptyset \text{ and } A \neq V_1\}$, one gets

$$-S_\bullet(\pi_e^{V_1})(s_1) = S_\bullet(\pi_e^{V_1})(-s_1) + \sum_{V_1 \cdot A \in V_1 \backslash \mathcal{U}_0(V_1)} S_+(\pi_{U_A}^{V_1}) S_\bullet(\pi_e^{U_A})((-1)^{i_A} s_1^{(V_1:U_A)}).$$

Hence, if $A_{k+1}, A_{k+2}, \dots, A_{k'} \in \mathcal{U}_0(V_1)$ denote representatives of the V_1 -orbits $V_1 A \subseteq \mathcal{U}_0(V_1)$ and we let $V_{k+1} := U_{A_{k+1}}, \dots, V_{k'} := U_{A_{k'}}$ then $V_i \leq V_1$ and $V \neq V_1$ for $i \geq k+1$. If we put $\varepsilon_{k+1} = \dots = \varepsilon_{k'} = 1$ and $s_{k+1} := (-1)^{i_{A_{k+1}}} s_1^{(V_1:V_{k+1})}, \dots, s_{k'} = (-1)^{i_{A_{k'}}} s_1^{(V_1:V_{k'})}$, then the polynomial

$$\xi_{(U; V_1, \dots, V_k; -1, \varepsilon_2, \dots, \varepsilon_k)}^G(s_1, \dots, s_k) := \xi_{(U; V_1, \dots, V_{k'}, 1, \varepsilon_2, \dots, \varepsilon_{k'})}^G(-s_1, s_2, \dots, s_{k'})$$

makes the statement of the lemma hold. We can conclude that if the lemma holds for every (n_1, n_2, n_3) with either $n_1 < m_1$ or $(n_1 = m_1, n_2 < m_2 \text{ and } n_3 \leq m_3)$, then it also holds for (m_1, m_2, m_3) .

Similarly, if $m_2 = 0$, but $m_3 > 0$, say $V_1 = V_2$, then we may use lemma 4.2 once more with $G = V_1$, $s = s_1$, and $t = s_2$ to conclude that

$$\begin{aligned} S_\bullet(\pi_e^{V_1})(s_1 + s_2) &= \sum_{V_1 \cdot A \in V_1 \setminus \mathcal{U}(V_1)} S_+(\pi_{U_A}^{V_1}) S_\bullet(\pi_e^{U_A})(s_1^{i_A} s_2^{i_{V_1-A}}) \\ &= S_\bullet(\pi_e^{V_1})(s_1) + S_\bullet(\pi_e^{V_1})(s_2) \\ &\quad + \sum_{V_1 \cdot A \in V_1 \setminus \mathcal{U}_0(V_1)} S_+(\pi_{U_A}^{V_1}) S_\bullet(\pi_e^{U_A})(s_1^{i_A} s_2^{i_{V_1-A}}), \end{aligned}$$

so with $V_{k+1}, \dots, V_{k'}$ as above, but with $\varepsilon_{k+1} = \dots = \varepsilon_{k'} = -1$ and with $s_{k+1} := s_1^{i_{A_{k+1}}} s_2^{i_{V_1-A_{k+1}}}, \dots, s_{k'} := s_1^{i_{A_{k'}}} s_2^{i_{V_1-A_{k'}}}$, the polynomial

$$\xi_{(U; V_1, \dots, V_k; \varepsilon_1, \dots, \varepsilon_k)}^G(s_1, \dots, s_k) := \xi_{(U; V_2, \dots, V_{k'}; 1, \varepsilon_2, \dots, \varepsilon_{k'})}^G(s_1 + s_2, s_3, \dots, s_{k'})$$

makes the statement of the lemma hold. We can conclude that if the lemma holds for every (n_1, n_2, n_3) with either $n_1 < m_1$ or $(n_1 = m_1, n_2 = m_2 = 0 \text{ and } n_3 < m_3)$, then it also holds for $(m_1, 0, m_3)$. The statement of the lemma now follows by induction first on m_1 , then on m_3 and finally on m_2 . $\square$

Lemma 4.5. *Let R be a ring. For some $k \in \mathbb{N}$ let $V_1, \dots, V_k \leq G$ be a sequence of subgroups of G and let $\varepsilon_1, \dots, \varepsilon_k \in \{\pm 1\}$. For every subgroup H of G one has, with the notation of lemma 4.4, that*

$$\sum_{i=1}^k \varepsilon_i |(G/V_i)^H| (s_i)^{(V_i:H)} = \sum'_{U \leq G} |(G/U)^H| (\xi_U(s_1, \dots, s_k))^{(U:H)}.$$

Proof. We use the same notation as in the above proof, and prove the lemma by induction on first m_1 , then on m_3 and finally on m_2 . Suppose that the lemma holds for every (n_1, n_2, n_3) with $n_1 < m_1$ or $n_3 < m_3$ or $(n_1 = m_1 \text{ and } n_3 = m_3 \text{ and } n_2 < m_2)$, and that we are given $(V_1, \dots, V_k; \varepsilon_1, \dots, \varepsilon_k)$ with $m_i = m_i(V_1, \dots, V_k; \varepsilon_1, \dots, \varepsilon_k)$. In the above proof we constructed $(W_1, \dots, W_{k'}; \delta_1, \dots, \delta_{k'})$ and $t_1, \dots, t_{k'} \in R$ such that $n_i = n_i(W_1, \dots, W_{k'}; \delta_1, \dots, \delta_{k'})$, $i = 1, 2, 3$ are on the form covered by the induction hypothesis, and such that:

$$\begin{aligned} \sum_{i=1}^k \varepsilon_i |(G/V_i)^H| s_i^{(V_i:H)} &= \sum_{i=1}^{k'} \delta_i |(G/W_i)^H| t_i^{(W_i:H)} \\ &= \sum'_{U \leq G} |(G/U)^H| (\xi_{(U; W_1, \dots, W_{k'}; \delta_1, \dots, \delta_{k'})}^G(t_1, \dots, t_{k'}))^{(U:H)} \\ &= \sum'_{U \leq G} |(G/U)^H| (\xi_{(U; V_1, \dots, V_k; \varepsilon_1, \dots, \varepsilon_k)}^G(s_1, \dots, s_k))^{(U:H)}. \end{aligned}$$

The other cases of the induction are similar. $\square$

Lemma 4.6. *For every G -Tambara functor S , for every two subgroups $V, W \leq G$ and for every $s, t \in S(G/e)$ one has the modified Mackey formula*

$$\begin{aligned} & S_+(\pi_V^G)S_\bullet(\pi_e^V)(s) \cdot S_+(\pi_W^G)S_\bullet(\pi_e^W)(t) \\ &= \sum_{VgW \in V \backslash G/W} S_+(\pi_{V \cap gWg^{-1}}^G)S_\bullet(\pi_e^{V \cap gWg^{-1}})(s^{(V:V \cap gWg^{-1})}) \cdot t^{(W:g^{-1}Vg \cap W)}. \end{aligned}$$

Proof. We use the diagram

$$\begin{array}{ccccc} G/e \amalg G/e & \xleftarrow{\text{pr}_1 \amalg \text{pr}_2} & G/e \times G/W \amalg G/V \times G/e & & \\ \pi_e^V \amalg \pi_e^W \downarrow & & \downarrow \pi_e^V \times \text{id} \amalg \text{id} \times \pi_e^W & & \\ G/G \amalg G/G & \xleftarrow{\pi_V^G \amalg \pi_W^G} & G/V \amalg G/W & \xleftarrow{\text{pr}_1 \amalg \text{pr}_2} & G/V \times G/W \amalg G/V \times G/W \\ \nabla \downarrow & & & & \nabla \downarrow \\ G/G & \longleftarrow & G/V \times G/W & \xlongequal{\quad} & G/V \times G/W, \end{array}$$

where ∇ is the fold map, the upper square is a pull-back and the lower rectangle is an exponential diagram. Concatenating with the diagram

$$\begin{array}{ccc} G/e \times G/W \amalg G/V \times G/e & \xleftarrow[\cong]{\alpha_1 \amalg \alpha_2} & \coprod_{gW \in G/W} G/e \amalg \coprod_{Vg \in V \backslash G} G/e \\ \pi_e^V \times \text{id} \amalg \text{id} \times \pi_e^W \downarrow & & \parallel \\ G/V \times G/W \amalg G/V \times G/W & & \coprod_{gW \in G/W} G/e \amalg \coprod_{Vg \in V \backslash G} G/e \\ \nabla \downarrow & & (\delta_1, \delta_2) \downarrow \\ G/V \times G/W & \xleftarrow[\cong]{\gamma} & \coprod_{VgW \in V \backslash G/W} G/(V \cap gWg^{-1}) \end{array}$$

with maps defined by

$$\begin{aligned} \alpha_1(gW, \sigma) &= (\sigma, \sigma gW), \\ \alpha_2(Vg, \sigma) &= (\sigma g^{-1}V, \sigma), \\ \delta_1(gW, \sigma) &= (VgW, \sigma(V \cap gWg^{-1})), \\ \delta_2(Vg, \sigma) &= (VgW, \sigma g^{-1}(V \cap gWg^{-1})) \quad \text{and} \\ \gamma(VgW, \sigma(V \cap gWg^{-1})) &= (\sigma V, \sigma gW), \end{aligned}$$

we get that

$$\begin{aligned} & S_+(\pi_V^G)S_\bullet(\pi_e^V)(s) \cdot S_+(\pi_W^G)S_\bullet(\pi_e^W)(t) \\ &= S[G/e \amalg G/e \leftarrow \coprod_{gW \in G/W} G/e \amalg \coprod_{Vg \in V \backslash G} G/e \rightarrow \coprod_{VgW \in V \backslash G/W} G/(V \cap gWg^{-1}) \rightarrow G/G](s, t) \\ &= \sum_{VgW \in V \backslash G/W} S_+(\pi_{V \cap gWg^{-1}}^G)S_\bullet(\pi_e^{V \cap gWg^{-1}})(s^{(V:V \cap gWg^{-1})}) \cdot t^{(W:g^{-1}Vg \cap W)}. \end{aligned}$$

□

Lemma 4.7 ([DS, Lemma 3.2.13]). *For every ring R , for every three subgroups $H, V, W \leq G$ and for every $s, t \in R$ one has the modified Mackey formula*

$$\begin{aligned} |(G/V)^H|_s^{(V:H)} |(G/W)^H|_t^{(W:H)} \\ = \sum_{VgW \in V \backslash G/W} |(G/V \cap gWg^{-1})^H|_{(s^{(V:V \cap gWg^{-1})} \cdot t^{(W:g^{-1}Vg \cap W)})}^{(V \cap gWg^{-1}:H)}. \end{aligned}$$

Proof of theorem 4.1. Let $G = V_1, V_2, \dots, V_k = U$ be a system of representatives of subgroups of G containing a conjugate of U . We define

$$s_U^G(a_{V_1}, b_{V_1}, \dots, a_{V_k}, b_{V_k}) := \xi_{(U; V_1, V_1, \dots, V_k, V_k; 1, \dots, 1)}^G(a_{V_1}, b_{V_1}, \dots, a_{V_k}, b_{V_k})$$

and

$$m_U^G(a_{V_1}, \dots, a_{V_k}) := \xi_{(U; V_1, \dots, V_k; -1, \dots, -1)}^G(a_{V_1}, \dots, a_{V_k}).$$

By the lemmas 4.4 and 4.5 these are integral polynomials with the desired properties. For example, for $U \leq G$ we have that

$$\begin{aligned} \phi_U(a) + \phi_U(b) &= \sum_{i=1}^k |(G/V_i)^U| (a_{V_i}^{(V_i:U)} + b_{V_i}^{(V_i:U)}) \\ &= \phi_U(\xi_{(U; V_1, V_1, \dots, V_k, V_k; 1, \dots, 1)}^G(a_{V_1}, b_{V_1}, \dots, a_{V_k}, b_{V_k})) \\ &= \phi_U(s_U^G(a_{V_1}, b_{V_1}, \dots, a_{V_k}, b_{V_k})). \end{aligned}$$

To construct $p_U = p_U^G$ we first choose a system $x_1, x_2, \dots, x_h$ of representatives of the G -orbits in

$$X := \coprod_{i,j=1}^k G/V_i \times G/V_j.$$

Next we put $W_r := G_{x_r}$ and

$$p_r = p_r(a_{V_1}, b_{V_1}, \dots, a_{V_k}, b_{V_k}) := a_i^{(V_i:W_r)} \cdot b_j^{(V_j:W_r)}$$

in case $x_r = (g_r V_i, g'_r V_j) \in G/V_i \times G/V_j \subseteq X$. Using these conventions, we define

$$p_U^G(a_{V_1}, b_{V_1}, \dots, a_{V_k}, b_{V_k}) := \xi_{(U; W_1, \dots, W_h; 1, \dots, 1)}^G(p_1, \dots, p_r).$$

Using the lemmas 4.6 and 4.7 we see that p_U has the desired properties. $\square$

5. FREE TAMBARA FUNCTORS

In this section we give a proof of theorem 3.6. On the way we shall give a combinatorial description of the Witt vectors of a polynomial G -ring, that is, a G -ring of the form $U^G(X, G/e)$ for a finite G -set X . Recall from 2.2 that the functor $R \mapsto R(G/e)$ from the category $[U^{fG}, \mathcal{E}ns]_0$ of fG -Tambara functors to the category of G -rings is an equivalence of categories, and that there are morphisms $(\mathcal{F}in^{fG})^{\text{op}} \subseteq (\mathcal{F}in^G)^{\text{op}} \xrightarrow{R} U^G$ of $\mathcal{Q}(G)$ -coloured theories. We let $F : \mathcal{E}ns^G \simeq [(\mathcal{F}in^G)^{\text{op}}, \mathcal{E}ns]_0 \rightarrow [U^G, \mathcal{E}ns]_0$ denote the left adjoint of the forgetful functor induced by the above composition of morphisms of coloured theories.

Definition 5.1. Given finite G -sets X and Y we let $\tilde{U}_+^G(X, Y) \subseteq \overline{U}^G(X, Y)$ denote those elements of the form $X \leftarrow A \rightarrow B \rightarrow Y$, where G acts freely on A , and we let $\tilde{U}^G(X, Y) \subseteq U^G(X, Y)$ denote the abelian subgroup generated by $\tilde{U}_+^G(X, Y)$. The composition

$$U^G(Y, Z) \times \tilde{U}^G(X, Y) \cup \tilde{U}^G(Y, Z) \times U^G(X, Y) \subseteq U^G(Y, Z) \times U^G(X, Y) \xrightarrow{\circ} U^G(X, Z)$$

factors through the inclusion $\tilde{U}^G(X, Z) \subseteq U^G(X, Z)$. We obtain a functor $\tilde{U}^G : \mathcal{F}in^G \rightarrow [U^G, \mathcal{E}ns]_0$ with $\tilde{U}^G(f : Y \rightarrow X) = \tilde{U}^G(R_f, -)$.

Lemma 5.2. *Given a G -Tambara functor S and a finite free G -set A , there is an isomorphism $\mathcal{E}ns^G(A, S(G/e)) \xrightarrow{\cong} S^*(A)$, which is natural in A .*

Proof. Choosing an isomorphism $\phi : A \xrightarrow{\cong} G/e \times A_0$ we obtain an isomorphism

$$\begin{aligned} \mathcal{E}ns^G(A, S(G/e)) &\xrightarrow{\phi^{-1*}} \mathcal{E}ns^G(G/e \times A_0, S(G/e)) \\ &\cong \mathcal{E}ns(A_0, S(G/e)) \cong S(G/e \times A_0) \xrightarrow{S^*(\phi)} S(A). \end{aligned}$$

This isomorphism is independent of the choice of ϕ . $\square$

Lemma 5.3. *Given a finite G -set X we have that $FX \cong \tilde{U}^G(X, -)$.*

Proof. For every G -Tambara functor S we shall construct a bijection

$$\mathcal{E}ns^G(X, S(G/e)) \cong [U^G, \mathcal{E}ns]_0(\tilde{U}^G(X, -), S).$$

Given $f : X \rightarrow S(G/e) \in \mathcal{E}ns^G(X, S(G/e))$ we let $\phi(f) \in [U^G, \mathcal{E}ns]_0(\tilde{U}^G(X, -), S)$ take $x = [X \xleftarrow{d} A \xrightarrow{b} B \xrightarrow{c} Y] \in \tilde{U}^G(X, Y)$ to $\phi(f)(x) \in S(Y)$ constructed as follows: by 5.2 we obtain an element $a \in S(A)$, and we let $\phi(f)(x) = S_+(c)S_\bullet(b)(a)$. Conversely, given $g \in [U^G, \mathcal{E}ns]_0(\tilde{U}^G(X, x), S)$, we construct $\psi(g) \in \mathcal{E}ns^G(X, S(G/e))$ by letting $\psi(g)(x) = [X \leftarrow G/e \rightarrow G/e \rightarrow G/e]$, where the map pointing left takes $e \in G$ to $x \in X$ and where the maps pointing right are identity maps. We leave it to the reader to check that ϕ and ψ are inverse bijections. $\square$

Corollary 5.4. *For every finite G -set X the functor $\tilde{U}^G(X, -) : U^G \rightarrow \mathcal{E}ns$ is isomorphic to $L_G U^G(X, G/e)$.*

Theorem 5.5. *Let X be a finite G -set and let $R = \tilde{U}^G(X, -)$. Then $R(G/e) = U^G(X, G/e)$, and the Teichmüller homomorphism*

$$\tau : \mathbb{W}_G(L_G R(G/e)) / \mathbb{I}_G(L_G R(G/e)) \rightarrow L_G R(G/G) = \tilde{U}^G(X, G/G)$$

is an isomorphism.

Proof of theorem B. Let X be a set with trivial action of G . The polynomial ring $\mathbb{Z}[X]$ has the universal property that determines the left adjoint of the forgetful functor from rings with an action of G to $\mathcal{E}ns^G$ up to isomorphism. Hence $\mathbb{Z}[X] \cong \tilde{U}^G(X, G/e) = U^G(X, G/e)$. By theorem 5.5 $\tilde{U}^G(X, G/G) = U^G(X, G/G)$ is isomorphic to $\mathbb{W}_G(\mathbb{Z}[X])$. $\square$

Proof of theorem 3.6. Let $R_0 = R(G/e)$. Given $\alpha = [W \leftarrow C \rightarrow D \rightarrow X] \in U^{fG}(W, X)$ we have an fG -Tambara map $\alpha^* : U^G(X, -) \rightarrow U^G(W, -)$ and we have the map $R(\alpha) : R(W) \rightarrow R(X)$. Hence we obtain maps

$$U^G(X, G/G) \times R(X) \leftarrow U^G(X, G/G) \times R(W) \rightarrow U^G(W, G/G) \times R(W).$$

Recall from lemma 2.4 that $L_G R(G/G)$ can be constructed by the coequalizer of the diagram

$$\coprod_{X, Y \in \text{ob } U^{fG}} U^G(X, G/G) \times R(Y) \rightrightarrows \coprod_{X \in \text{ob } U^{fG}} U^G(X, G/G) \times R(X),$$

induced by the above maps. We shall construct a map

$$\rho : L_G R(G/G) \rightarrow \mathbb{W}_G(R_0)/\mathbb{I}_G(R_0)$$

by specifying explicit maps $\rho_X : U^G(X, G/G) \times R(X) \rightarrow \mathbb{W}_G(R_0)/\mathbb{I}_G(R_0)$. Given $\underline{r} \in R(X)$, we have an fG -Tambara morphism $\text{ev}_{\underline{r}} : U^{fG}(X, -) \rightarrow R$. In particular we have a G -homomorphism $\text{ev}_{\underline{r}}(G/e) : U^G(X, G/e) \rightarrow R(G/e) = R_0$. By theorem 5.5 we get an induced ring-homomorphism

$$U^G(X, G/G) \cong \mathbb{W}_G(U^G(X, G/e))/\mathbb{I}_G(U^G(X, G/e)) \rightarrow \mathbb{W}_G(R_0)/\mathbb{I}_G(R_0),$$

and hence by adjunction we obtain a map

$$\rho_X : U^G(X, G/G) \times R(X) \rightarrow \mathbb{W}_G(R_0)/\mathbb{I}_G(R_0).$$

We need to check that these ρ_X induce a map on the coequalizer $L_G R(G/G)$ of the above coequalizer diagram, that is, for α as above we need to show that the diagram

$$\begin{array}{ccc} U^G(X, G/G) \times R(W) & \xrightarrow{\alpha^* \times \text{id}} & U^G(W, G/G) \times R(W) \\ \text{id} \times R(\alpha) \downarrow & & \rho_W \downarrow \\ U^G(X, G/G) \times R(X) & \xrightarrow{\rho_X} & \mathbb{W}_G(R_0)/\mathbb{I}_G(R_0) \end{array}$$

commutes. In order to do this we note that the diagram

$$\begin{array}{ccc} \mathbb{W}_G(U^G(X, G/e)) & \xrightarrow{t} & U^G(X, G/G) \\ \alpha^* \downarrow & & \alpha^* \downarrow \\ \mathbb{W}_G(U^G(W, G/e)) & \xrightarrow{t} & U^G(W, G/G) \end{array}$$

commutes, and therefore it will suffice to show that the diagram

$$\begin{array}{ccc} \mathbb{W}_G(U^G(X, G/e)) \times R(W) & \xrightarrow{\alpha^* \times \text{id}} & \mathbb{W}_G(U^G(W, G/e)) \times R(W) \\ \text{id} \times R(\alpha) \downarrow & & \downarrow \\ \mathbb{W}_G(U^G(X, G/e)) \times R(X) & \longrightarrow & \mathbb{W}_G(R_0) \end{array}$$

commutes, where the arrows without labels are constructed using the homomorphisms $\mathbb{W}_G(\text{ev}_{\underline{r}}(G/e))$ for $\underline{r}$ an element of either $R(X)$ or $R(W)$. Using diagonal

inclusions of the form

$$\mathbb{W}_G(T) \times Z \rightarrow \mathbb{W}_G(T) \times \prod'_{U \leq G} Z \approx \prod'_{U \leq G} (T \times Z)$$

we see that it suffices to show note that the diagram

$$\begin{array}{ccc} \prod'_{U \leq G} (U^G(X, G/e) \times R(W)) & \xrightarrow{\prod'_{U \leq G} (\alpha^* \times \text{id})} & \prod'_{U \leq G} (U^G(W, G/e) \times R(W)) \\ \prod'_{U \leq G} (\text{id} \times R(\alpha)) \downarrow & & \downarrow \\ \prod'_{U \leq G} (U^G(X, G/e) \times R(X)) & \longrightarrow & \prod'_{U \leq G} R(G/e) \approx \mathbb{W}_G(R_0) \end{array}$$

commutes. This finishes the construction of $\rho : L_G R(G/G) \rightarrow \mathbb{W}_G(R_0)/\mathbb{I}_G(R_0)$. We leave it to the reader to check that ρ and τ are inverse bijections. For this it might be helpful to note that

$$\begin{aligned} & \sum'_{U \leq G} ([G/e \xleftarrow{=} G/e \xrightarrow{\pi_e^U} G/U \rightarrow G/G] \circ [Y \leftarrow A_U \rightarrow B_U \rightarrow G/e]) \\ &= [\prod'_{U \leq G} G/e \xleftarrow{=} \prod'_{U \leq G} G/e \rightarrow \prod'_{U \leq G} G/U \rightarrow G/G] \circ [Y \leftarrow \prod'_{U \leq G} A_U \rightarrow \prod'_{U \leq G} B_U \rightarrow \prod'_{U \leq G} G/e]. \end{aligned}$$

□

In order to begin the proof of theorem 5.5 we need to introduce filtrations of both sides.

Definition 5.6. Let R be a ring and let $U \leq G$ be a subgroup of G . We let $I_U \subseteq \mathbb{W}_G(R)$ denote the ideal generated by those $a = (a_K)'_{K \leq G} \in \mathbb{W}_G(R)$ satisfying that $a_K \neq 0$ implies that $K \lesssim U$. We let $\tilde{I}_U \subseteq I_U$ denote the subideal $\tilde{I}_U = \sum_{V \subsetneq U} I_V \subseteq I_U$.

Definition 5.7. Given a G -sets X , we let $J_U^+ = J_U(X, G/G) \subseteq \tilde{U}^G(X, G/G)$ denote the subset of elements of the form $[X \leftarrow A \rightarrow B \rightarrow G/G] \in U_+^G(X, G/G) \subseteq U^G(X, G/G)$, where $B^K = \emptyset$ when $U \lesssim K$ and $U \neq K$. We let $J_U \subseteq U^G(X, G/G)$ denote the ideal generated by J_U^+ and we let $\tilde{J}_U \subseteq J_U$ denote the subideal $\tilde{J}_U = \sum_{V \subsetneq U} J_V \subseteq J_U$.

Lemma 5.8. *In the situation of the above definition the following holds:*

- (i) *Any element in J_U is of the form $x - y$ for $x, y \in J_U^+$.*
- (ii) *Every element x in the image of the map $J_U^+ \rightarrow J_U/\tilde{J}_U$ is of the form*

$$x = [X \xleftarrow{d} G/e \times A \xrightarrow{\pi_e^U \times f} G/U \times B \xrightarrow{q} G/G] + \tilde{J}_U,$$

with $f : A \rightarrow B$ a map of (nonequivariant) sets and d a G -map, where q is the composition $G/U \times B \xrightarrow{\text{pr}} G/U \xrightarrow{\pi_U^G} G/G$.

- (iii) *If*

$$x = [X \xleftarrow{d} G/e \times A \xrightarrow{\pi_e^U \times f} G/U \times B \xrightarrow{q} G/G] + \tilde{J}_U,$$

and

$$x' = [X \xleftarrow{d'} G/e \times A' \xrightarrow{\pi_e^U \times f'} G/U \times B' \xrightarrow{q'} G/G] + \tilde{J}_U,$$

then $x = x'$ if and only if there exist bijections $\alpha : A \rightarrow A'$ and $\beta : B \rightarrow B'$ and for every $a \in A$ there exists $g_a \in N_G(U)$ such that firstly $d'(e, \alpha a) = d(g_a, a)$, and secondly, if $a_1, a_2 \in A$ satisfy that $f(a_1) = f(a_2)$, then $g_{a_1}U = g_{a_2}U$.

Proof. A straight forward verification yields that multiplication in $\tilde{U}_+^G(X, G/G)$ induces a map $J_U^+ \times \tilde{U}_+^G(X, G/G) \rightarrow J_U^+$ and that J_U^+ is closed under sum. It follows that J_U is the abelian subgroup of $U^G(X, G/G)$ generated by J_U^+ . The statement (i) is a direct consequence of this. For (ii) we note that for every element

$$r = [X \xleftarrow{d} D \xrightarrow{c} E \xrightarrow{t} G/G]$$

in $\tilde{U}_+^G(X, G/G)$, we have a decomposition $E \cong \coprod'_{K \leq G} E_K$, where $E_K \cong G/K \times B_K$ for some B_K . This decomposition induces an isomorphism

$$\tilde{U}^G(X, G/G) \cong \bigoplus'_{K \leq G} J_K / \tilde{J}_K$$

of abelian groups. Given an element x of the form

$$x = [X \xleftarrow{d} D \xrightarrow{c} G/U \times B \xrightarrow{q} G/G] + \tilde{J}_U,$$

we can choose a G -bijection of the form $c^{-1}(G/U \times \{b\}) \cong G/e \times A_b$ for every $b \in B$. It follows that x is represented by an element of the form

$$x = [X \xleftarrow{d} G/e \times A \xrightarrow{p \times f} G/U \times B \xrightarrow{q} G/G] + \tilde{J}_U.$$

We leave the straight forward verification of part (iii) to the reader. $\square$

Lemma 5.9. *Let U be a subgroup of G and let $a = (a_U)'_{U \leq G} \in I_U$ have*

$$a_U = [X \xleftarrow{d} A \times G/e \xrightarrow{f \times 1} B \times G/e \xrightarrow{\text{pr}} G/e].$$

Then

$$\tau(a) \equiv [X \xleftarrow{d} A \times G/e \xrightarrow{f \times \pi_e^U} B \times G/U \rightarrow G/G] \mod \tilde{J}_U.$$

Proof. The lemma follows from the diagram

$$\begin{array}{ccccccc} X & \xleftarrow{d} & A \times G/e & \longleftarrow & W & \longleftarrow & G/e \times A \\ & & f \times \text{id} \downarrow & & \downarrow & & \downarrow \\ G/e & \xleftarrow{\text{pr}} & G/e \times B & \longleftarrow & Z & \longleftarrow & G/e \times B \\ \pi_e^U \downarrow & & & & \downarrow & & \downarrow \\ G/U & \longleftarrow & \coprod_{gU \in G/U} \text{map}(gU, B) & \equiv & Y & \longleftarrow & G/U \times B, \end{array}$$

where the lower rectangle is an exponential diagram and the squares are pull-backs. We use that the map $G/U \times B \rightarrow Y$ which takes (gU, b) to the constant map $gU \rightarrow B$ with value b is an isomorphism on G/U -parts and that $Y^H = \emptyset$ for $U \lesssim H$ and $U \neq H$. $\square$

Corollary 5.10. *Let X be a G -set and let $R = \tilde{U}^G(X, -)$. The map*

$$\tau : \mathbb{W}_G(U^G(X, G/e)) \rightarrow L_G R(G/G) = \tilde{U}^G(X, G/G)$$

satisfies that $\tau(I_U) \subseteq J_U$ and that $\tau(\tilde{I}_U) \subseteq \tilde{J}_U$.

Proposition 5.11. *Let X be a G -set, let $R = \tilde{U}^G(X, -)$ and let $R_0 = R(G/e)$. For every $U \leq G$ the map $\tau : \mathbb{W}_G(R_0)/\mathbb{I}_G(R_0) \rightarrow L_G R(G/G) = \tilde{U}^G(X, G/G)$ induces an isomorphism $\tau_U : (\mathbb{I}_G(R_0) + I_U)/(\mathbb{I}_G(R_0) + \tilde{I}_U) \rightarrow J_U/\tilde{J}_U$.*

Proof. Let $x \in I_U$ have $x_U = [X \xleftarrow{d} A \times G/e \xrightarrow{f \times 1} B \times G/e \xrightarrow{p} G/e]$. Then by lemma 5.9 $\tau(x) \equiv [X \xleftarrow{d} A \times G/e \xrightarrow{f \times p} B \times G/U \xrightarrow{q} G/G] \bmod \tilde{J}_U$, with the notation introduced there, and it follows from lemma 5.8 that τ_U is onto. On the other hand, to prove injectivity, we pick $x_1, x_2 \in I_U$ with $\tau(x_1) \equiv \tau(x_2) \bmod \tilde{J}_U$. Suppose that $x_{i,U}$ has the form

$$x_{i,U} = [Z \xleftarrow{d_i} A_i \times G/e \xrightarrow{f_i \times 1} B_i \times G/e \xrightarrow{p} G/e]$$

for $i = 1, 2$. Let

$$y_i = [Z \xleftarrow{d_i} A_i \times G/e \xrightarrow{f_i \times \pi_e^U} B \times G/U \xrightarrow{q} G/G]$$

for $i = 1, 2$. Then by lemma 5.9 $y_i \equiv \tau(x_i) \bmod \tilde{J}_U$ for $i = 1, 2$. It follows from lemma 5.8 that there exist bijections $\alpha : A_1 \rightarrow A_2$ and $\beta : B_1 \rightarrow B_2$ and for every $a \in A_1$ there exists $g_a \in N_G(U)$ such that firstly $d_1(\alpha a) = g_a d(a)$ and secondly, if $a_1, a_2 \in A$ satisfy that $f(a_1) = f(a_2)$, then $g_{a_1} U = g_{a_2} U$. Given $a \in A_i$, let $z_{i,a} \in \tilde{U}^G(X, G/e)$ denote the element $[X \xleftarrow{d_{i,a}} G/e \xrightarrow{p} G/e \xrightarrow{p} G/e]$, where $d_{i,a}(e) = d_i(a, e)$. Then $z_{2,\alpha(a)} = g_a z_{1,a}$ and $x_{i,U} = \sum_{b \in B_i} (\prod_{a \in f_i^{-1}(b)} z_{i,a})$ for $i = 1, 2$, where an empty product is 1 and an empty sum is 0. We can conclude that $x_{1,U} - x_{2,U} \in \mathbb{I}_G(R)$, and hence $x_1 - x_2 \in \mathbb{I}_G(R) + \tilde{I}_U$. In the general case $\tau(x_1 - x'_1) \equiv \tau(x_2 - x'_2) \bmod \tilde{J}_U$ we easily obtain that $x_1 - x'_1 \equiv x_2 - x'_2 \bmod \mathbb{I}_G(R) + \tilde{I}_U$ by collecting the positive terms. $\square$

Proof of theorem 5.5. We first note that $\tilde{I}_V = \sum_{U \subsetneq V} I_U \cong \operatorname{colim}_{U \subsetneq V} I_U \subseteq I_V$ and that $\tilde{J}_V = \sum_{U \subsetneq V} J_U \cong \operatorname{colim}_{U \subsetneq V} J_U \subseteq J_V$. The result now follows by induction on the cardinality of V using the above proposition and the five lemma on the following map of short exact sequences:

$$\begin{array}{ccccc} \tilde{\mathbb{I}}_G(R(G/e)) + I_V & \longrightarrow & \mathbb{I}_G(R(G/e)) + I_V & \longrightarrow & \frac{(\mathbb{I}_G(R(G/e)) + I_V)}{(\mathbb{I}_G(R(G/e)) + \tilde{I}_V)} \\ \downarrow & & \downarrow & & \downarrow \\ \tilde{J}_V & \longrightarrow & J_V & \longrightarrow & J_V/\tilde{J}_V. \end{array}$$

$\square$

6. THE WITT TAMBARA-FUNCTOR

In this section we finally prove theorem C. Given a subgroup $H \leq G$ and an H -set X , we can construct a $G \times H$ -set $G/e \times X$, where G acts by multiplication on the left on G/e , and where $h \cdot (g, x) := (gh^{-1}, hx)$. We let $\text{ind}_H^G X$ denote the G -set $H \backslash (G/e \times X)$.

Lemma 6.1. *Let H be a subgroup of G . The functor $\text{ind}_H^G : \mathcal{F}in^H \rightarrow \mathcal{F}in^G$ induces functors $\text{ind}_H^G : U^H \rightarrow U^G$, and $\text{ind}_{fH}^{fG} : U^{fH} \rightarrow U^{fG}$.*

Proof. Since the functor $\text{ind}_H^G : \mathcal{F}in^H \rightarrow \mathcal{F}in^G$ preserves pull-back diagrams and exponential diagrams it induces a functor $\text{ind}_H^G : U^H \rightarrow U^G$ that takes $X \leftarrow A \rightarrow B \rightarrow Y$ to $\text{ind}_H^G X \leftarrow \text{ind}_H^G A \rightarrow \text{ind}_H^G B \rightarrow \text{ind}_H^G Y$. $\square$

Given a G -Tambara functor S we can construct an H -Tambara functor $\text{res}_H^G S = S \circ \text{ind}_H^G$. Similarly, given an fG -Tambara functor R we can construct an fH -Tambara functor $\text{res}_{fH}^{fG} S = S \circ \text{ind}_{fH}^{fG}$.

Theorem 6.2. *Given an fG -Tambara functor R , the Teichmüller homomorphism*

$$\tau : \mathbb{W}_H(\text{res}_{fH}^{fG} R(H/e)) / \mathbb{I}_H(\text{res}_{fH}^{fG} R(H/e)) \rightarrow \text{res}_H^G L_G(H/H)$$

is an isomorphism.

Proof of theorem C. If we consider $R_0 = R(G/e)$ as an H -ring, then

$$\mathbb{W}_H(\text{res}_{fH}^{fG}(H/e)) / \mathbb{I}_H(\text{res}_{fH}^{fG}(H/e)) \cong \mathbb{W}_H(R_0) / \mathbb{I}_H(R_0),$$

and $\text{res}_H^G L_G(H/H) = L_G R(G/H)$. Combining these observations with 6.2 we obtain the statement of theorem C. $\square$

Lemma 6.3. *Let H be a subgroup of G . The forgetful functor $i^* : \mathcal{F}in^G \rightarrow \mathcal{F}in^H$ which takes a G -set Y to the same set considered as an H -set induces functors $i^* : U^G \rightarrow U^H$ and $i_f^* : U^{fG} \rightarrow U^{fH}$.*

Lemma 6.4. *Let H be a subgroup of G . The functor $i^* : U^G \rightarrow U^H$ is left adjoint to the functor $\text{ind}_H^G : U^H \rightarrow U^G$ and the functor $i_f^* : U^{fG} \rightarrow U^{fH}$ is left adjoint to the functor $\text{ind}_{fH}^{fG} : U^{fH} \rightarrow U^{fG}$.*

Proof. We prove only the first part of the lemma. Given $X \leftarrow A \rightarrow B \rightarrow G \times_H Y$ in $U^G(X, G \times_H Y)$ we construct an element in $U^H(i^* X, Y)$ by the following diagram where the two squares furthest to the right are pull-back squares:

$$\begin{array}{ccccccc} i^* X & \longleftarrow & A_H & \longrightarrow & B_H & \longrightarrow & H \times_H Y \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ X & \longleftarrow & A & \longrightarrow & B & \longrightarrow & G \times_H Y \end{array}$$

Conversely, given $i^* X \leftarrow E \rightarrow F \rightarrow Y$ in $U^H(i^* X, Y)$ we construct the element $X \leftarrow G \times_H E \rightarrow G \times_H F \rightarrow G \times_H Y$ in $U^G(X, G \times_H Y)$. Here the arrow pointing to the left is the composite $G \times_H E \rightarrow G \times_H i^* X \rightarrow X$. We leave it to the reader to check that the maps are inverse bijections in an adjunction. $\square$

We have the following commutative diagram of categories:

$$\begin{array}{ccc} Uf^G & \xrightarrow{i_f^*} & Uf^H \\ \downarrow & & \downarrow \\ U^G & \xrightarrow{i^*} & U^H, \end{array}$$

where the vertical functors are the natural inclusions. Since ind_H^G is right adjoint to i^* , res_H^G is left adjoint to $[i^*, \mathcal{E}ns]_0$ (see for example [Sch, Proposition 16.6.3].) Similarly res_{fH}^{fG} is left adjoint to $[i_f^*, \mathcal{E}ns]_0$. From the commutative diagram of functor categories

$$\begin{array}{ccc} [Uf^G, \mathcal{E}ns]_0 & \xleftarrow{[i_f^*, \mathcal{E}ns]_0} & [Uf^H, \mathcal{E}ns]_0 \\ \uparrow & & \uparrow \\ [U^G, \mathcal{E}ns]_0 & \xleftarrow{[i^*, \mathcal{E}ns]_0} & [U^H, \mathcal{E}ns]_0, \end{array}$$

where the vertical maps are the forgetful functors induced by the inclusions $Uf^G \subseteq U^G$ and $Uf^H \subseteq U^H$ we can conclude that there is a natural isomorphism $\text{res}_H^G L_G \cong L_H \text{res}_{fH}^{fG}$.

Proof of theorem 6.2. By theorem 3.6 we have an isomorphism

$$\mathbb{W}_H(\text{res}_{fH}^{fG} R(H/e)) / \mathbb{I}_H(\text{res}_{fH}^{fG} R(H/e)) \xrightarrow{\tau} L_H \text{res}_{fH}^{fG} R(H/H) \cong \text{res}_H^G L_G R(H/H).$$

□

REFERENCES

- [AR] J. Adamek, J. Rosicky, *Locally Presentable and Accessible Categories*. London Mathematical Society Lecture Note Series 189, (1994).
- [Be] P. Berthelot, *Cohomologie cristalline des schemas de caracteristique $p > 0$* . Lecture Notes in Mathematics. 407. (1974).
- [BV] J. Boardman, R. Vogt, *Homotopy invariant algebraic structures on topological spaces*. Springer-Verlag, Berlin, (1973), Lecture Notes in Mathematics, Vol. 347.
- [Bo] F. Borceux, *Handbook of categorical algebra 2*. Cambridge University Press, Cambridge (1994), Categories and Structures.
- [Br] M. Brun, *Equivariant stable homotopy and Tambara functors*. In preparation.
- [DS] A. Dress, C. Siebeneicher *The Burnside Ring of Profinite Groups and the Witt Vector Construction*, Advances in Math. 70 (1988).
- [DS2] A. Dress, C. Siebeneicher *The Burnside ring of the infinite cyclic group and its relations to the necklace algebra, λ -rings, and the universal ring of Witt vectors*. Adv. Math. 78, No.1, 1-41 (1989).
- [Gra] J. Graham, *Generalised Witt vectors*. Adv. Math. 99, No.2, 248-263 (1993).
- [Li] H. Lindner, *A remark on Mackey-functors*. Manuscr. Math. 18, 273-278 (1976)
- [MR] N. Metropolis, G-C. Rota, *Witt vectors and the algebra of necklaces*. Adv. Math. 50, 95-125 (1983)
- [Re] C. Rezk, *Every homotopy theory of simplicial algebras admits a proper model*. Topology Appl. 119, No.1, 65-94 (2002).
- [Sch] H. Schubert, *Categories*. Springer-Verlag (1972).
- [SW] H. Schmid, E. Witt *Unverzweigte abelsche Körper vom Exponenten p^n über einem algebraischen Funktionenkörper der Charakteristik p* . J. Reine Angew. Math. 176, 168-173 (1936).

- [Ta] D. Tambara *On multiplicative transfer*. Commun. Algebra 21, No.4, 1393-1420 (1993).
- [Wi] E. Witt, *Zyklische Körper und Algebren der Charakteristik p vom Grad p^n . Struktur diskret bewerteter perfekter Körper mit vollkommenem Restklassenkörper der Charakteristik p* . J. Reine Angew. Math. 176, 126-140 (1936).

E-mail address: `brun@mathematik.uni-osnabrueck.de`

FB MATHEMATIK/INFORMATIK, UNIVERSITÄT OSNABRÜCK, 49069 OSNABRÜCK, GERMANY